

IDENTITIES INVOLVING q -GENOCCHI NUMBERS AND POLYNOMIALS

SERKAN ARACI, MEHMET ACIKGOZ, HASSAN JOLANY, AND YUAN HE

ABSTRACT. In this paper, we focus on the q -Genocchi numbers and polynomials. We shall introduce new identities of the q -Genocchi numbers and polynomials by using the fermionic p -adic integral on $\mathbb{Z}_p$ which are very important in the study of Frobenius-Genocchi numbers and polynomials. Also, we give Cauchy-integral formula for the q -Genocchi polynomials and moreover by using measure theory on p -adic integral we derive the distribution formula q -Genocchi polynomials. Finally, we present a new definition of q -Zeta-type function by using Mellin transformation which is the interpolation function of the q -Genocchi polynomials at negative integers.

1. PRELIMINARIES

Recently, *Kim* and *Lee* have given some properties for the q -Euler numbers and polynomials in [7]. Actually, this type numbers and polynomials and their q -extensions or variously generalizations have been studied in several different ways for a long time (for details, see [1-21]).

By using p -adic q -integral on $\mathbb{Z}_p$, *Kim* defined many new generating functions of the q -Bernoulli polynomials, q -Euler polynomials and q -Genocchi polynomials in his arithmetic works (for details, see [6-19]). The works of *Kim* have been benefited for further works of many mathematicians in Analytic numbers theory. Actually, we motivated from his inspiring works to write this paper.

Assume that p be a fixed odd prime number. Throughout this work, we require the definitions of the some notations such that let $\mathbb{Q}_p$ be the field p -adic rational numbers and let $\mathbb{C}_p$ be the completion of algebraic closure of $\mathbb{Q}_p$. That is,

$$\mathbb{Q}_p = \left\{ x = \sum_{n=-k}^{\infty} a_n p^n : 0 \leq a_n \leq p-1 \right\}.$$

Then $\mathbb{Z}_p$ is integral domain which is defined by

$$\mathbb{Z}_p = \left\{ x = \sum_{n=0}^{\infty} a_n p^n : 0 \leq a_n \leq p-1 \right\}$$

or

$$\mathbb{Z}_p = \left\{ x \in \mathbb{Q}_p : |x|_p \leq 1 \right\}.$$

2000 *Mathematics Subject Classification.* Primary 05A10, 11B65; Secondary 11B68, 11B73.

Key words and phrases. Genocchi numbers and polynomials, q -Genocchi numbers and polynomials, p -adic q -integral on $\mathbb{Z}_p$, Mellin transformation, q -Zeta function.

We assume that $q \in \mathbb{C}_p$ with $|1 - q|_p < 1$ as an indeterminate. The p -adic absolute value $|\cdot|_p$, is normally given by

$$|x|_p = p^{-r}$$

where $x = p^r \frac{s}{t}$ with $(p, s) = (p, t) = (s, t) = 1$ and $r \in \mathbb{Q}$.

The q -extension of x with the display notation of $[x]_q$ is introduced by

$$[x]_q = \frac{1 - q^x}{1 - q}.$$

We want to note that $\lim_{q \rightarrow 1} [x]_q = x$ (see[1-22]).

Also, we use notation $\mathbb{N}^*$ for the union of zero and natural numbers.

We consider that η is a uniformly differentiable function at a point $a \in \mathbb{Z}_p$, if the difference quotient

$$\Phi_\eta(x, y) = \frac{\eta(x) - \eta(y)}{x - y},$$

have a limit $\eta'(a)$ as $(x, y) \rightarrow (a, a)$ and denote this by $\eta \in UD(\mathbb{Z}_p)$. Then, for $\eta \in UD(\mathbb{Z}_p)$, we can discuss the following

$$\frac{1}{[p^n]_q} \sum_{0 \leq \xi < p^n} \eta(\xi) q^\xi = \sum_{0 \leq \xi < p^n} \eta(\xi) \mu_q(\xi + p^n \mathbb{Z}_p),$$

which represents as a p -adic q -analogue of Riemann sums for η . The integral of η on $\mathbb{Z}_p$ will be defined as the limit ($n \rightarrow \infty$) of these sums, when it exists. The p -adic q -integral of function $\eta \in UD(\mathbb{Z}_p)$ is defined by T. Kim in [6], [10], [15] by

$$(1.1) \quad I_q(\eta) = \int_{\mathbb{Z}_p} \eta(\xi) d\mu_q(\xi) = \lim_{n \rightarrow \infty} \frac{1}{[p^n]_q} \sum_{\xi=0}^{p^n-1} \eta(\xi) q^\xi.$$

The bosonic integral is considered as a bosonic limit $q \rightarrow 1$, $I_1(\eta) = \lim_{q \rightarrow 1} I_q(\eta)$. Similarly, the fermionic p -adic integral on $\mathbb{Z}_p$ is introduced by T. Kim as follows:

$$(1.2) \quad I_{-q}(\eta) = \int_{\mathbb{Z}_p} \eta(\xi) d\mu_{-q}(\xi)$$

(for more details, see [16-18]). Obviously that

$$(1.3) \quad \lim_{q \rightarrow 1} I_{-q}(\eta) = I_{-1}(\eta) = \int_{\mathbb{Z}_p} \eta(\xi) d\mu_{-1}(\xi) = \lim_{n \rightarrow \infty} \sum_{\xi=0}^{p^n-1} \eta(\xi) (-1)^\xi.$$

From (1.3), it is well-known as the useful property for the fermionic p -adic q -integral on $\mathbb{Z}_p$:

$$(1.4) \quad I_{-1}(\eta_1) + I_{-1}(\eta) = 2\eta(0),$$

where $\eta_1(x) = \eta(x+1)$ (for details, see[2-4, 11-14, 16-22]).

The q -Genocchi polynomials can be introduced as follows:

$$(1.5) \quad G_{n,q}(x) = n \int_{\mathbb{Z}_p} q^\xi (x + \xi)^{n-1} d\mu_{-1}(\xi).$$

From (1.5), we have

$$G_{n,q}(x) = \sum_{l=0}^n \binom{n}{l} x^l G_{n-l,q}$$

where $G_{n,q}(0) := G_{n,q}$ are called q -Genocchi numbers. Then, q -Genocchi numbers can be given by

$$G_{0,q} = 0 \text{ and } q(G_q + 1)^n + G_{n,q} = \begin{cases} 2, & \text{if } n = 1 \\ 0, & \text{if } n \geq 1 \end{cases}$$

with the usual convention about replacing $(G_q)^n$ by $G_{n,q}$ is used (for details, see [4]).

Our objective in the present paper is to derive not only new but also novel and interesting properties of the q -Genocchi numbers and polynomials. Our applications for the q -Genocchi polynomials seem to be useful in mathematics for engineerings (on this subject, see [23]).

2. SOME PROPERTIES ON THE q -GENOCCHI NUMBERS AND POLYNOMIALS

Let $\eta(x) = q^x e^{t(x+\xi)}$. Then, by using (1.4), we see that

$$t \int_{\mathbb{Z}_p} q^\xi e^{t(x+\xi)} d\mu_{-1}(\xi) = \frac{2t}{qe^t + 1} e^{xt}.$$

From the last equality and (1.5), we easily derive the following generating function of the q -Genocchi polynomials:

$$(2.1) \quad t \int_{\mathbb{Z}_p} q^\xi e^{t(x+\xi)} d\mu_{-1}(\xi) = \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!} = \frac{2t}{qe^t + 1} e^{xt}, \quad |\log q + t| < \pi.$$

Substituting $x \rightarrow x + y$ into (2.1), then we write

$$\begin{aligned} \sum_{n=0}^{\infty} G_{n,q}(x+y) \frac{t^n}{n!} &= \frac{2t}{qe^t + 1} e^{(x+y)t} \\ &= \left(\sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!} \right) \left(\sum_{n=0}^{\infty} y^n \frac{t^n}{n!} \right) \\ &= \sum_{n=0}^{\infty} \left(\sum_{k=0}^n \binom{n}{k} G_{k,q}(x) y^{n-k} \right) \frac{t^n}{n!} \end{aligned}$$

From the above, we easily express the following theorem:

Theorem 1. *The following holds:*

$$(2.2) \quad G_{n,q}(x+y) = \sum_{k=0}^n \binom{n}{k} G_{k,q}(x) y^{n-k}.$$

By (2.2), we consider the following

$$G_{n,q}(x+y) = \frac{2}{[2]_q} ny^{n-1} + \sum_{k=2}^n \binom{n}{k} G_{k,q}(x) y^{n-k}.$$

From this, it follows that

$$G_{n,q}(x+y) - \frac{2}{[2]_q} ny^{n-1} = \sum_{k=2}^n \binom{n}{k} G_{k,q}(x) y^{n-k}$$

can be derived and so we reach the following theorem:

Theorem 2. For $n \in \mathbb{N}^*$, one has

$$(2.3) \quad \sum_{k=0}^n \frac{\binom{n}{k}}{(k+2)(k+1)} G_{k+2,q}(x) y^{n-k} \\ = \frac{G_{n+2,q}(x+y) - \frac{2}{[2]_q} (n+2) y^{n+1}}{(n+2)(n+1)}.$$

Replacing y by $-y$ into (2.3), then we get

$$(2.4) \quad \frac{G_{n+2,q}(x-y) - \frac{2}{[2]_q} (n+2) (-1)^{n+1} y^{n+1}}{(n+2)(n+1)} \\ = \sum_{k=0}^n \frac{\binom{n}{k} (-1)^{n-k}}{(k+2)(k+1)} G_{k+2,q}(x) y^{n-k}.$$

By (2.4), it follows that

$$(2.5) \quad \sum_{k=0}^n \frac{\binom{n}{k} (-1)^k}{(k+2)(k+1)} G_{k+2,q}(x) y^{n-k} \\ = \frac{(-1)^n G_{n+2,q}(x-y) + \frac{2}{[2]_q} (n+2) y^{n+1}}{(n+1)(n+2)}.$$

Therefore, from the expressions of (2.3) and (2.5), we procure the following theorem:

Theorem 3. The following holds true:

$$(2.6) \quad \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} \frac{\binom{n}{2k}}{(k+1)(2k+1)} G_{2k+2,q}(x) y^{n-2k} \\ = \frac{(-1)^n G_{n+2,q}(x-y) + G_{n+2,q}(x+y)}{(n+1)(n+2)}$$

where $[\cdot]$ is Gauss' symbol.

By (2.4), we have the following identity

$$(2.7) \quad \sum_{k=2}^n \frac{\binom{n}{k} (-1)^k}{k(k-1)} G_{k,q}(x) y^{n-k} \\ = \frac{(-1)^n G_{n,q}(x-y) + \frac{2}{[2]_q} n y^{n-1}}{n(n-1)}.$$

By (2.4), (2.5) and (2.7), then we have the following theorem:

Theorem 4. For $n \in \mathbb{N}^*$, we get

$$(2.8) \quad \frac{(-1)^n G_{n+2,q}(x-y) + G_{n+2,q}(x+y)}{(n+2)(n+1)} \\ = \sum_{k=0}^{\lfloor \frac{n+1}{2} \rfloor} \frac{\binom{n}{2k}}{(k+1)(2k+1)} G_{2k+2,q}(x) y^{n-2k}.$$

Taking $y = 1$ into (2.3), it follows that

$$(2.9) \quad q \sum_{k=0}^n \frac{\binom{n}{k}}{(k+2)(k+1)} G_{k+2,q}(x) = \frac{qG_{n+2,q}(x+1)}{(n+1)(n+2)} - \frac{2q}{[2]_q(n+1)}.$$

We need the following for sequel of this paper:

$$2e^{tx} = \frac{1}{t} \left(q \frac{2t}{qe^t + 1} e^{(x+1)t} + \frac{2t}{qe^t + 1} e^{xt} \right).$$

From the above, we easily develop the following:

$$(2.10) \quad qG_{n+1,q}(x+1) + G_{n+1,q}(x) = (n+1)2x^n.$$

By (2.9) and (2.10), we state the following theorem:

Theorem 5. *The following holds:*

$$(2.11) \quad \sum_{k=0}^n \frac{\binom{n}{k}}{(k+2)(k+1)} G_{k+2,q}(x) = \frac{2x^{n+1}}{qn+q} - \frac{G_{n+2,q}(x)}{(qn+q)(n+2)} - \frac{2}{[2]_q(n+1)}.$$

Thanks to equality of $\lim_{q \rightarrow 1} G_{n,q}(x) = G_{n,1}(x) := G_n(x)$, where $G_n(x)$ are known as Genocchi polynomials which is given via the exponential generating function, as follows:

$$\sum_{n=0}^{\infty} G_n(x) \frac{t^n}{n!} = \frac{2t}{e^t + 1} e^{xt} \quad (|t| < \pi),$$

(see [1-4, 12, 13, 21]). From the above, as $q \rightarrow 1$ in (2.11), we discover the following corollary:

Corollary 1. *The following*

$$\begin{aligned} & \sum_{k=0}^n \frac{\binom{n}{k}}{(k+2)(k+1)} G_{k+2}(x) \\ &= \frac{2x^{n+1}}{n+1} - \frac{G_{n+2}(x)}{(n+1)(n+2)} - \frac{1}{n+1} \end{aligned}$$

is true.

Let us take $y = 1$ and $n \rightarrow 2n$ into (2.6), becomes

$$(2.12) \quad \begin{aligned} & \sum_{k=0}^n \frac{\binom{2n}{2k}}{(k+1)(2k+1)} G_{2k+2,q}(x) \\ &= \frac{G_{2n+2,q}(x-1) + G_{2n+2,q}(x+1)}{(2n+1)(2n+2)} \\ &= \frac{\frac{1}{q}(qG_{2n+2,q}(x+1) + G_{2n+2,q}(x)) + qG_{2n+2,q}(x) + G_{2n+2,q}(x-1)}{(2n+1)(2n+2)} \\ & \quad - \frac{G_{2n+2,q}(x)}{q(2n+1)(2n+2)} - \frac{qG_{2n+2,q}(x)}{(2n+1)(2n+2)} \\ &= \frac{2(n+2)x^{n+1}}{(2n+1)(2n+2)} + \frac{2(n+2)(x-1)^{n+1}}{(2n+1)(2n+2)} - \frac{G_{2n+2,q}(x)}{q(2n+1)(2n+2)} - \frac{qG_{2n+2,q}(x)}{(2n+1)(2n+2)} \end{aligned}$$

After these applications, we conclude with the following theorem:

Theorem 6. *The following identity*

$$(2.13) \quad \sum_{k=0}^n \frac{\binom{2n}{2k}}{(k+1)(2k+1)} G_{2k+2,q}(x) \\ = \frac{(n+2)x^{n+1}}{(2n+1)(n+1)} + \frac{(n+2)(x-1)^{n+1}}{(2n+1)(n+1)} - \frac{G_{2n+2,q}(x)}{q(2n+1)(2n+2)} - \frac{qG_{2n+2,q}(x)}{(2n+1)(2n+2)}$$

is true.

Now, we analyse as $q \rightarrow 1$ for the equation (2.13) and so we readily state the following corollary which seems to be interesting property for the Genocchi polynomials.

Corollary 2. *The following equality holds:*

$$(2.14) \quad \sum_{k=0}^n \frac{\binom{2n}{2k}}{(k+1)(2k+1)} G_{2k+2}(x) \\ = \frac{2(n+2)x^{n+1}}{(2n+1)(2n+2)} + \frac{2(n+2)(x-1)^{n+1}}{(2n+1)(2n+2)} - \frac{G_{2n+2}(x)}{(2n+1)(2n+2)} - \frac{G_{2n+2}(x)}{(2n+1)(2n+2)}.$$

Substituting $n \rightarrow 2n+1$ and $y=1$ into (2.8), we compute

$$\begin{aligned} & \sum_{k=0}^n \frac{\binom{2n+1}{2k}}{(k+1)(2k+1)} G_{2k+2,q}(x) \\ &= \frac{G_{2n+3,q}(x+1) - G_{2n+3,q}(x-1)}{(2n+3)(2n+2)} \\ &= \frac{\frac{1}{q}(qG_{2n+3,q}(x+1) + G_{2n+3,q}(x)) - (G_{2n+3,q}(x) + G_{2n+3,q}(x-1))}{(2n+3)(2n+2)} \\ &+ \left(\frac{q-1}{q}\right) \frac{G_{2n+3,q}(x)}{(2n+3)(2n+2)} \\ &= \frac{x^{2n+2}}{q(n+1)} - \frac{(x-1)^{2n+2}}{(n+1)} + \left(\frac{q-1}{q}\right) \frac{G_{2n+3,q}(x)}{(2n+3)(2n+2)}. \end{aligned}$$

Therefore, we obtain the following theorem:

Theorem 7. *The following equality holds:*

$$\begin{aligned} & \sum_{k=0}^n \frac{\binom{2n+1}{2k}}{(k+1)(2k+1)} G_{2k+2,q}(x) \\ &= \frac{x^{2n+2}}{q(n+1)} - \frac{(x-1)^{2n+2}}{(n+1)} + \left(\frac{q-1}{q}\right) \frac{G_{2n+3,q}(x)}{(2n+3)(n+1)}. \end{aligned}$$

As $q \rightarrow 1$ in the above theorem, then we easily derive the following corollary:

Corollary 3. *For $n \in \mathbb{N}^*$, then we have*

$$\begin{aligned} & \sum_{k=0}^n \frac{\binom{2n+1}{2k}}{(k+1)(2k+1)} G_{2k+2,q}(x) \\ &= \frac{2x^{2n+2}}{(2n+2)} - \frac{2(x-1)^{2n+2}}{(2n+2)}. \end{aligned}$$

3. CONCLUSION

In this final section, we recall the generating function of the q -Genocchi polynomials, as follows:

$$(3.1) \quad \mathcal{F}_q(x, t) = \frac{2t}{qe^t + 1} e^{xt} = \sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!}.$$

Using the definition of the k -th derivative as $\frac{d^k}{dt^k}$ to (3.1), then we easily see that

$$(3.2) \quad \frac{d^k}{dt^k} \left(\frac{2t}{qe^t + 1} e^{xt} \right) = \frac{d^k}{dt^k} \left(\sum_{n=0}^{\infty} G_{n,q}(x) \frac{t^n}{n!} \right).$$

By applying $\lim_{t \rightarrow 0}$ on the both sides in (3.2), then we conclude with the following theorem:

Theorem 8. *The following identity*

$$(3.3) \quad G_{k,q}(x) = \lim_{t \rightarrow 0} \left[\frac{d^k}{dt^k} \left(\frac{2t}{qe^t + 1} e^{xt} \right) \right]$$

is true.

We now consider Cauchy-integral formula of the q -Genocchi polynomials which is a vital and important in complex analysis, is an important statement about line integrals for holomorphic functions in the complex plane. So, by using the equation of (3.3), we can state the following theorem:

Theorem 9. *The following Cauchy-integral holds true:*

$$G_{n,q}(x) = \frac{n!}{2\pi i} \int_C \mathcal{F}_q(x, t) \frac{dt}{t^{n+1}}$$

where C is a loop which starts at $-\infty$, encircles the origin once in the positive direction, and the returns $-\infty$.

Distribution formula for the q -Genocchi polynomials is important to study regarding p -adic Measure theory. That is,

$$\begin{aligned} \int_{\mathbb{Z}_p} q^y (x+y)^n d\mu_{-1}(y) &= \lim_{n \rightarrow \infty} \sum_{\xi=0}^{dp^n-1} (-1)^\xi (x+\xi)^n q^\xi \\ &= d^n \sum_{a=0}^{d-1} (-1)^a q^a \left(\lim_{n \rightarrow \infty} \sum_{\xi=0}^{p^n-1} (-1)^\xi \left(\frac{x+a}{d} + \xi \right)^n q^{d\xi} \right) \\ &= d^n \sum_{a=0}^{d-1} (-1)^a q^a \frac{G_{n+1,q}\left(\frac{x+a}{d}\right)}{n+1}. \end{aligned}$$

After the above applications, we procure the following theorem.

Theorem 10. *For $n \in \mathbb{N}^*$, then we have*

$$G_{n,q}(dx) = d^{n-1} \sum_{a=0}^{d-1} (-1)^a q^a G_{n,q}\left(x + \frac{a}{d}\right).$$

By utilizing from the definition of the geometric series in (3.1), we easily see that

$$\sum_{m=0}^{\infty} G_{m,q}(x) \frac{t^m}{m!} = \sum_{m=0}^{\infty} \left(2(m+1) \sum_{n=0}^{\infty} (-1)^n q^n n^m \right) \frac{t^{m+1}}{(m+1)!},$$

by comparing the coefficients on the both sides, then we have, for $m \in \mathbb{N}$

$$(3.4) \quad \frac{G_{m+1,q}(x)}{m+1} = [2]_q \sum_{n=1}^{\infty} (-1)^n q^n n^m.$$

Now also, we develop the following applications by using Mellin transformation to the generating function of the q -Genocchi polynomials: For $s \in \mathbb{C}$ and $\Re(s) > 1$,

$$(3.5) \quad \begin{aligned} \zeta(s, x : q) &= \frac{1}{\Gamma(s)} \int_0^{\infty} t^{s-2} \{-\mathcal{F}_q(x, -t)\} dt \\ &= 2 \sum_{n=0}^{\infty} (-1)^n q^n \left(\int_0^{\infty} t^{s-1} e^{-nt} dt \right) \\ &= 2 \sum_{n=1}^{\infty} \frac{(-1)^n q^n}{n^s} \end{aligned}$$

Thus, we can state the definition of the q -Zeta-type function as follows:

$$(3.6) \quad \zeta(s, x : q) = 2 \sum_{n=1}^{\infty} \frac{(-1)^n q^n}{n^s}.$$

As $q \rightarrow 1$ in (3.6), turns into

$$\lim_{q \rightarrow 1} \zeta(s, x : q) = \zeta(s, x : 1) := \zeta(s, x) = 2 \sum_{n=1}^{\infty} \frac{(-1)^n}{n^s}$$

which is well-known as Euler-Zeta function (see [6]). By (3.4) and (3.6), we get

$$\zeta(-n, x : q) = \frac{G_{n+1,q}(x)}{n+1}.$$

REFERENCES

- [1] S. Araci, D. Erdal, and J. J. Seo, A study on the fermionic p -adic q -integral representation on $\mathbb{Z}_p$ associated with weighted q -Bernstein and q -Genocchi polynomials, *Abstract and Applied Analysis*, Volume **2011**, Article ID 649248, 10 pages.
- [2] S. Araci, J. J. Seo, and D. Erdal, New Construction weighted (h, q) -Genocchi numbers and polynomials related to Zeta type function, *Discrete Dynamics in Nature and Society*, Volume **2011**, Article ID 487490, 7 pages.
- [3] S. Araci, M. Acikgoz, and Feng Qi, On the q -Genocchi numbers and polynomials with weight zero and their applications, Available at <http://arxiv.org/abs/1202.2643>
- [4] H. Jolany and H. Sharifi, Some results for the Apostol-Genocchi polynomials of higher order, *Bulletin of Malaysian Mathematical Sciences Society* (Article in press).
- [5] Yuan He and Chunping Wang, Some Formulae of Products of the Apostol-Bernoulli and Apostol-Euler Polynomials, *Discrete Dynamics in Nature and Society*, vol. **2012**, Article ID 927953, 11 pages.
- [6] T. Kim, Euler numbers and polynomials associated with Euler Zeta functions, *Abstract and Applied Analysis*, Volume **2008**, Article ID 581582, 11 pages.
- [7] T. Kim and S.-H. Lee, Some Properties on the q -Euler Numbers and Polynomials, *Abstract and Applied Analysis*, vol. **2012**, Article ID 284826, 9 pages, 2012.
- [8] D. Kim, T. Kim, S.-H. Lee, D. V. Dolgy, S.-H. Rim, Some new identities on the Bernoulli numbers and polynomials, *Discrete Dynamics in Nature and Society*, Volume **2011**, Article ID 856132, 11 pages.

- [9] T. Kim, J. Choi, and Y. H. Kim, Some identities on the q -Bernoulli numbers and polynomials with weight 0, *Abstract And Applied Analysis*, Volume **2011**, Article ID 361484, 8 pages.
- [10] T. Kim, On a q -analogue of the p -adic log gamma functions related integrals, *J. Number Theory*, 76 (**1999**) no. 2, 320-329.
- [11] T. Kim, and J. Choi, On the q -Euler numbers and polynomials with weight 0, *Abstract and Applied Analysis*, Volume **2012**, Article ID 795304, 7 pages..
- [12] T. Kim, On the q -extension of Euler and Genocchi numbers, *J. Math. Anal. Appl.* 326 (**2007**) 1458-1465.
- [13] T. Kim, On the multiple q -Genocchi and Euler numbers, *Russian J. Math. Phys.* 15 (4) (**2008**) 481-486.
- [14] T. Kim, On the weighted q -Bernoulli numbers and polynomials, *Advanced Studies in Contemporary Mathematics* 21(**2011**), no.2, p. 207-215.
- [15] T. Kim, q -Volkenborn integration, *Russ. J. Math. phys.* 9 (**2002**) 288-299.
- [16] T. Kim, q -Euler numbers and polynomials associated with p -adic q -integrals, *J. Nonlinear Math. Phys.*, 14 (**2007**), no. 1, 15–27.
- [17] T. Kim, New approach to q -Euler polynomials of higher order, *Russ. J. Math. Phys.*, 17 (**2010**), no. 2, 218–225.
- [18] T. Kim, Some identities on the q -Euler polynomials of higher order and q -Stirling numbers by the fermionic p -adic integral on $\mathbb{Z}_p$, *Russ. J. Math. Phys.*, 16 (**2009**), no.4,484–491.
- [19] D. Kim, T. Kim, J. Choi, and Y. H. Kim, Identities involving q -Bernoulli and q -Euler numbers, *Abstract and Applied Analysis*, Volume **2012** (2012), Article ID 674210, 10 pages.
- [20] C. S. Ryoo, A note on the weighted q -Euler numbers and polynomials, *Advan. Stud. Contemp. Math.* 21(**2011**), 47-54.
- [21] I. N. Cangul, V. Kurt, H. Ozden, and Y. Simsek, On the higher-order w - q -genocchi numbers, *Advanced Studies in Contemporary Mathematics* 19 (**2009**), no. 1, pp. 39–57.
- [22] M. Acikgoz, S. Araci and I. N. Cangul, A note on the modified q -Bernstein polynomials for functions of several variables and their reflections on q -Volkenborn integration, *Applied Mathematics and Computation*, vol. 218 (**2011**), no. 3, pp. 707–712.
- [23] L. C. Andrews, Special Functions of Mathematics for Engineerings, *SPIE Press*, 1992, pages 479.

**University of Gaziantep, Faculty of Science and Arts, Department of Mathematics,
27310 Gaziantep, TURKEY**

E-mail address: mtsrkn@hotmail.com; saraci88@yahoo.com.tr

**University of Gaziantep, Faculty of Science and Arts, Department of Mathematics,
27310 Gaziantep, TURKEY**

E-mail address: acikgoz@gantep.edu.tr

**School of Mathematics, Statistics and Computer Science, University of Tehran,
Iran**

E-mail address: hassan.jolany@khayam.ut.ac.ir; hassan.jolany@mail.com

**Department of Mathematics, Kunming University of Science and Technology,
Kunming, Yunnan 650500, People' Republic of China**

E-mail address: hyyhe@yahoo.com.cn